arXiv:1110.3213v1 [cond-mat.quant-gas] 14 Oct 2011

Incompressible Quantum Glass State of Bosons in Two Dimensions

Yancheng Wang,[1] Wenan Guo,[1, *] and Anders W. Sandvik[2, †]

[1]*Department of Physics, Beijing Normal University, Beijing 100875, China*
[2]*Department of Physics, Boston University, 590 Commonwealth Avenue, Boston, Massachusetts 02215, USA*

(Dated: October 17, 2011)

We study the quantum glass state intervening between the conventional superfluid and Mott-insulator states of lattice bosons with random potentials. Its properties at temperature $T = 0$ are controlled by rare large regions of superfluid surrounded by Mott insulator. These regions make the state gapless although it is insulating. Contrary to the commonly accepted theory of this state in two dimensions, we show here that a vanishing gap does not necessarily imply nonzero compressibility. Using quantum Monte Carlo simulations of the Bose-Hubbard model and a percolation theory, we show that the compressibility κ at average filling $\rho = 1$ follows the form $\kappa \sim \exp(-b/T^{\alpha})$ with $\alpha < 1$. The system is, thus, incompressible at $T = 0$ and should be classified as a Mott glass.

PACS numbers: 67.85.-d, 67.85.Hj, 67.10.F, 64.70.Tg

Interacting lattice bosons have two distinct types of ground states in the absence of disorder; the superfluid (SF) and the Mott-insulator (MI). In the standard prototypical Bose-Hubbard (BH) model with repulsive on-site interactions, an MI state has an integer number of particles per site and there is a gap to states with added or removed particles. The gapless SF can have any filling fraction. These phases and the quantum phase transitions between them are well understood [1–5] and have also been realized experimentally with ultracold atoms in optical lattices [6, 7]. If disorder in the form of random site potentials is introduced in the BH model a third kind of state appears—an insulating but gapless quantum glass (QG). This state has been the subject of numerous studies over the past two decades [1–3, 8–20], but many of its properties are still not well understood. In one dimension, two typses of QG ground states are known; the compressible Bose glass (BG) [1], and the incompressible Mott glass (MG) [19], with the latter appearing at commensurate filling fractions only [20]. In two and higher dimensions it has been believed that the QG is always of the compressible BG type [2, 14]. QG states are also accessible in optical lattices [21, 22].

We here discuss site-disordered bosons at temperature $T > 0$. Our arguments and conclusions will mostly be generic, but for definiteness, to study systems using unbiased quantum Monte Carlo (QMC) simulations, we consider the two-dimensional (2D) BH Hamiltonian

$$H = -t \sum_{\langle ij \rangle} (b_i^+ b_j + b_j^+ b_i) + \frac{U}{2} \sum_{i=1}^{N} n_i(n_i - 1) + \sum_{i=1}^{N} \epsilon_i n_i, \quad (1)$$

where $\langle ij \rangle$ denotes nearest-neighbor sites on the square lattice, b_i^+ (b_i) are boson creation (destruction) operators, $n_i = b_i^+ b_i$ site occupation numbers, and ϵ_i random potentials uniformly distributed in the range $[-\Lambda - \mu, \Lambda - \mu]$ around the average chemical potential μ.

Previous studies of the site-disordered BH model have focused on its $T = 0$ phase diagram and, in particular, the long-standing question of whether or not a QG state always intervenes between the SF and MI phases in the presence of disorder [2, 3, 8–18]. Such questions arise in many classical and quantum many-body problems, with the QG corresponding generically to a Griffiths phase in which statistically rare large regions of some phase A inside another phase B lead to singularities characterizing a new type of disordered phase (Griffiths phase) not present in the absence of disorder [23–25]. In the case of the BH model with bounded disorder (which is necessary to retain an MI state), the Griffiths argument states the following [2, 3]: In the MI, once the width 2Λ of the disorder distribution exceeds the Mott gap Δ_M, there can be arbitrarily large regions of SF inside the MI. These SF "puddles" are not immediately, until Λ exceeds some larger critical value, percolating through the lattice, however, so the state is insulating. The QG is still fundamentally different from the MI because it is gapless. This scenario was recently refined and named the "theorem of inclusion" [14].

While, according to the theorem, the MI–QG boundary is dictated exactly by the Mott gap of the clean MI [3, 14], it has been pointed out that this boundary cannot be extracted based on QMC simulations, because the rare regions leading to the vanishing gap are in practice not generated on the relatively small lattices to which the calculations are limited [3, 14, 18]. Indeed, some earlier numerical studies [8, 10] had partially missed the QG phase because its signatures at $T = 0$ are difficult to distinguish from those of the MI. Renormalization group (RG) studies as well had pointed to the possibility of a direct MI–SF transition [11, 12].

The purpose of our work presented here is to clarify the consequences of gaplessness of the 2D QG—in particular, to show that the absence of a gap does not necessarily imply a compressible state. We will argue that, in contrast to the commonly accepted theory of 2D site-disordered bosons [2, 3, 13, 14, 18], the compressibility κ of the QG can vanish when $T \to 0$. For the model (1) at filling



FIG. 1: (Color online) Disorder-averaged compressibility versus inverse temperature for systems with $\rho = 1$, $U = 22$, and different values of the width Λ of the disorder distribution. The lattice size is $L = 32$ in all cases. The curves are fits to the exponential form (2) with $\alpha = 1$ for $\Lambda = 0, 3$ (MI state) and $\alpha = 0.77$ and 0.53 for $\Lambda = 6$ and 7, respectively (MG state). For $\Lambda = 9$ (SF state) the line is horizontal.

fraction $\rho = 1$ we find the form

$$\kappa \sim \exp(-b/T^{\alpha}), \qquad (2)$$

where $\alpha < 1$. In the MI phase the presence of a gap Δ and a slowly varying density of states at the gap edge implies the above form with $\alpha = 1$ and $b = \Delta$. The incompressibility of the QG at $T = 0$ puts it in the MG class, which in site-disordered systems was up until now only found in one dimension [19].

We next present QMC results demonstrating (2) and then discuss some aspects of the density of states and how the anomalous exponential compressibility follows from an SF quantum percolation scenario.

QMC calculations—We have used the stochastic series expansion QMC method with directed-loop updates [26] to study the Hamiltonian (1) with the chemical potential μ adjusted to give an average filling fraction $\rho = \langle n \rangle / N = 1$ (typically to accuracy 10^{-5} or better) in averages over hundreds of realizations of the random potentials. To speed up the simulations we restrict the site occupation to $n_i \leq 2$, which for large U only has minor unimportant consequences (which we have checked explicitly by some calculations with $n_i \leq 3$). We fix $U/t = 22$. According to previous work [18] (with which our results agree well) the system is then an MI for $\Lambda < 4.2$, an SF for $\Lambda > 7.8$, and a QG in the window between these points (and also in a separate region for $\Lambda > 70$ that we do not study here). For inverse temperatures $\beta = t/T$ up to 8, we eliminate finite-size effects by using $L \times L$ lattices with L up to 32. Results for $L = 16$ and $L = 32$ agree within error bars.

As shown in Fig. 1, the behavior in the QG (MG) phase follows the form (2) with $\alpha \approx 0.77$ for $\Lambda = 6$ and $\alpha \approx 0.53$ for $\Lambda = 7$. For points inside the MI phase ($\Lambda = 0$ and 3) the behavior agrees with the conventional $\alpha = 1$ exponential as expected (and also discussed further below). Inside the SF phase, exemplified in the figure by results for $\Lambda = 9$, the compressibility converges rapidly to a non-zero value as T is lowered, as also expected.



FIG. 2: (Color online) Compressibility distribution for $L = 16$ systems at different inverse temperatures. The total number of instances of the random potentials was 1200 for each β.

An important question is whether the low-T κ distribution develops a stretched tail, which would make it difficult to obtain reliable disorder averages based on a relatively small number of samples. This problem at $T = 0$ was stressed in [3, 14]. Fig. 2 shows histograms for $L = 16$ systems. It is clear that the distribution becomes increasingly skewed as the temperature is lowered, with a thin high-κ tail developing. The peak moves rapidly toward lower values, however, and the tail is not sufficiently long or fat to cause serious problems with properly converging the average κ. We have confirmed that successive binning of the samples into increasingly large groups leads to convergent error bars (which in Fig. 1 are smaller than the plot symbols in most cases). It is likely that the tails will become very significant at even lower temperatures, but with the whole distribution shifting toward lower values, so that the average $\kappa \to 0$ as $T \to 0$ according to the anomalous exponential form (2).

Compressibility and density of states—According to Eq. (2) the QG is incompressible at $T = 0$, i.e., it is an MG, in contrast to the long-held belief that it should be a compressible BG [2, 3, 13, 14]. The gaplessness in the thermodynamic limit follows from the presence of rare large regions of SF surrounded by MI. An infinite SF domain is of course compressible and leads to a vanishing gap of the system as a whole. The probability of a given site belonging to an infinitely large SF region is vanishingly small, however, and the compressibility normalized by the system volume in the standard way can therefore vanish when $T \to 0$ despite the absence of a gap.

We will next consider the basic relationships between the density of states and the compressibility, to demonstrate more formally why an MG with $\kappa \to 0$ is plausible. To this end, we write the compressibility as

$$\kappa = \frac{1}{TN} \frac{\sum_n \int d\epsilon \mathrm{e}^{-\epsilon/T} \rho_n(\epsilon)(n - \langle n \rangle)^2}{\sum_n \int d\epsilon \mathrm{e}^{-\epsilon/T} \rho_n(\epsilon)}, \qquad (3)$$

where $\rho_n(\epsilon)$ is the density of states with n particles.

Consider first the MI state of a clean system. There is a gap Δ to adding a particle. In a simplified picture, valid for large U, the added particle should disperse as a free boson, i.e., with the 2D energy-momentum relation $\epsilon_q = \Delta + 4t - 2t[\cos(q_x) + \cos(q_y)]$. This gives an almost flat density of states at the band edges and a van Hove singularity at the center. At low temperatures the singularity should not matter and the compressibility should be governed by the density of states at the lower band edge. We can therefore consider a simplified flat density of states, which gives the same low-T behavior (up to unimportant T and N independent factors).

For definiteness, we again consider a system with average density $\rho = 1$. There are N single-particle states, for both added and removed particles, and the energies in these sectors are the same due to the emergent [2] particle-hole symmetry. We distribute the states uniformly within a band of width W above the gap Δ. Thus, the density of these states is $\rho_{N\pm1}(\epsilon) = N/W$ for $\epsilon \in [\Delta, \Delta + W]$ and 0 otherwise. The compressibility at low T, where we only need to consider $n = N$ in the denominator of (3), is then $\kappa = (2/W)\mathrm{e}^{-\Delta/T}$. The pure exponential form agrees with our $\Lambda = 0$ and $\Lambda = 3$ QMC results in Fig. 1. For $\Lambda > 0$ Δ should correspond to an average gap, smaller than the gap of the clean system.

Note that, in a clean system with perfect particle-hole symmetry, the thermodynamic gap Δ controlling κ as above should be equal to half the Mott gap Δ_M. The $\Lambda = 0$ fit in Fig. 1 gives $\Delta = 4.3(1)$, which indeed agrees well with $\Delta_M \approx 8.34$ extracted from the difference in chemical potential between the upper and lower edges of the $\rho = 1$ Mott lobe at $U = 22$.

When the system is gapless it is no longer true that the ground state dominates the partition function at low temperatures. Consider the $n = N \pm 1$ sectors. When $N \to \infty$ we can neglect the ground state contribution to the partition function in (3) if $\rho_{N\pm1}(\epsilon \to 0) > 0$ or if $\rho_{N\pm1}(\epsilon)$ vanishes sufficiently slowly when $\epsilon \to 0$. Since $(n - N)^2 = 1$ we get a compressibility contribution $\propto 1/NT$, independently of the exact form of the density of states. Thus, the contribution from the single-particle sector vanishes for any $T > 0$ as $N \to \infty$.

In the gapless case we can of course not neglect the sectors with more than one added or removed particle, because those sectors should also be gapless, at least when the deviation from the ground state occupation number $\delta_n \ll N$. In order for the total compressibility to be size independent (non-zero when $N \to \infty$) in the limit $T \to 0$, it is crucial, however, that there are enough low-energy states also in sectors with δ_n up to order $\mathcal{O}(\sqrt{N})$. This is required in order to cancel the factor $1/N$ in the normalization of κ in (3).

In the standard picture of the QG, a significant density of low-energy states with $\mathcal{O}(\sqrt{N})$ fluctuations seems unlikely, because the probability of a given site belonging to an SF region (which is where the particle number can fluctuate) decreases exponentially with the size of the region. Finite SF domains should have gaps between states in different particle number sectors, and, thus, become incompressible as $T \to 0$. These SF domain gaps will lead to a very low density of low-energy states for the sectors with n deviating significantly from N. The scenario $\kappa \to 0$ as $T \to 0$, suggested by our numerical results, therefore appears very plausible.

Percolation scenario—The form (2) can be understood heuristically using percolation theory: Consider an ensemble of SF domains below the percolation point. At $T = 1/m^a$ all domains of sizes $s < m$ are effectively in their ground states and do not contribute to the compressibility. The exponent a depends on the low-energy level spectrum of the SF domains, which may be affected by the domain shapes and boundaries. Domains with $s > m$ should contribute essentially independently of T and s. The probability of a given site belonging to a domain with $s > m$ is $\propto \exp(-bm^c)$, where c is known in classical percolation theory [27] but may be different in a quantum system. In terms of the unknown exponents a and c the compressibility due to SF domains is $\kappa \propto \exp(-bT^{-c/a})$, which is Eq. (2) with $\alpha = c/a$.

One might worry about the validity of this picture when the SF domains are not completely independent, being weakly coupled through the surrounding MI. However, since the SF is not percolating through the system, the level structure should not be affected beyond a broadening of the individual domain levels on a much smaller scale than the level spacing. It remains to be understood how α can depend on the BH model parameters.

Cross-over behavior—We have not yet carried out a systematic study of the BH model on a dense grid of disorder strengths Λ, but from the results at two different MG points in Fig. 1 we know that the exponent α depends on the system parameters. There should always be a contribution $\propto \exp(-\Delta/T)$ to the compressibility from the MI background (with Δ an average gap), but this becomes negligible at low T when $\alpha < 1$. Upon approaching the MG–MI transition, b in (2) should diverge. The relative MI contribution then grows, so that κ crosses over to the form with $\alpha = 1$. At the $T = 0$ MG–SF transition one expects $\kappa \sim (\Lambda_c - \Lambda)^{\nu(2-z)}$ with $z = 2$ (as in the generic clean MI–SF transition) [2]. As the transition is approached from the MG, $\alpha \to 0$ would ensure that κ can evolve smoothly for any $T > 0$ between the MG form and the constant κ exactly at the transition (with a discontinuity developing as $T \to 0$). Both the transitions out of the MG should be studied in more detail to clarify the exact nature of the cross-overs. The behavior at the MG–SF transition, in particular, should shed light on the mechanism of quantum percolation of the SF puddles.

In a previous QMC study of a (2 + 1)-dimensional classical model with a glassy phase, mimicking the 2D quantum BH model (1), it was claimed that the $T = 0$ compressibility at integer filling is non-singular at the SF transition [13] and nonzero in the glass phase. However, if $\alpha \to 0$ at this transition (and also in other possible cross-over scenarios), simulations carried out at low but non-zero temperature in the MG in close vicinity of the SF boundary may produce results that appear to converge to a non-zero value, even though $\kappa \to 0$ eventually as $T \to 0$ according to (2) with a small α. In [13] $\kappa > 0$ was indeed seen only in simulations very close to the SF.

Conclusions and discussion—We have presented plausible arguments and unbiased QMC results demonstrating an incompressible MG phase in the BH model. Previous arguments for a generically compressible BG, starting with the seminal work by Fisher *et al.* [2] and continuing up to more recent studies [14, 18], have never, to our knowledge, been backed up by complete proofs taking into account the probability of $\mathcal{O}(\sqrt{N})$ particle-number fluctuations. The outlines of proofs presented in Refs. [2] and [14] seem to neglect the fact that the normalization by $1/N$ in Eq. (3) makes $\kappa \to 0$ possible even though there are SF domains with arbitrarily small gaps.

Our key physical argument is that a finite SF domain surrounded by an MI in the BH model becomes incompressible as $T \to 0$. This is clear for a single SF domain in a large MI, where the ground state has a definite number of particles and there must be a gap to states with added or removed particles. The gap decreases with increasing size of the SF domain. In the QG state, with many SF domains surrounded by MI, as long as the SF cannot percolate through the system (which would bring it into the SF state) it should effectively consist of isolated SF puddles with fixed particle numbers at $T = 0$. If this picture is correct the anomalous exponential form (2) follows, independently of dimensionality.

While we have carried out substantial QMC calculations only at density $\rho = 1$ (in part because it is much more difficult to locate QG states at significantly lower fixed average density), there is no apparent reason, based on the percolation scenario, why QG states of the BH model with non-integer average density should be qualitatively different. In principle it is possible, however, that the emergent particle-hole symmetry [2] at integer ρ in the clean system survives in the presence of disorder (as it can do in one dimension [19, 20]) and makes these systems special. Indeed, according to a recent weak-disorder RG calculation [16], 2D systems with integer filling have a different fixed point that is not accessible within that RG (and nothing could be said about the compressibility). Previous works [2, 14] found no distinction between integer and noninteger filling. Our results at $\rho = 1$ may then correspond to this unknwon fixed point, and the possibility remains of a compressible BG existing at noninteger filling. This would also imply that the percolation picture does not apply at noninteger fillings.

The percolation scenario can, in fact, break down in one dimension, where, according to strong-disorder RG [20], the SF domains at incommensurate filling grow as the energy scale is lowered but there is no long-range superfluidity because of one dimensional localization due to the disorder. This mechanism is very particular to one dimension, however, and does not directly carry over to higher dimensions. Whether or not commensurate fillings are special in 2D has to be tested with numerical simulations at non-integer filling.

We note that the same type of anomalous exponential behavior (2) was previously observed in the magnetic susceptibility of a quantum spin system with a glass phase similar to the MG [28]. In that case $\alpha = 1/2$ was found in all cases studied, in contrast to the continuously evolving α we have found here in the BH model. One reason for the different α could be that $z = 1$ in the spin system.

Acknowledgments—We would like to thank Claudio Chamon, Matthew Fisher, Philip Phillips, and Anatoli Polkovnikov for useful discussions. This work was supported by the NSFC under Grant No. 11175018 and NCET-08-0053 (WG) and by the NSF under Grant No. DMR-1104708 (AWS). WG would like to thank the Condensed Matter Theory Visitors program at Boston University. AWS gratefully acknowledges support from NCET-08-0053 for a visit to Beijing Normal University.

---

[*] e-mail: waguo@bnu.edu.cn
[†] e-mail: sandvik@bu.edu

[1] T. Giamarchi and H. J. Schulz, Phys. Rev. B **37**, 325 (1988).
[2] M. P. A. Fisher, P. B. Weichman, G. Grinstein, and D. S. Fisher, Phys. Rev. B **40**, 546 (1989).
[3] J. K. Freericks and H. Monien, Phys. Rev. B **53**, 2691 (1996).
[4] G. G. Batrouni, V. Rousseau, R. T. Scalettar, M. Rigol, A. Muramatsu, P. J. H. Denteneer, and M. Troyer, Phys. Rev. Lett. **89**, 117203 (2002).
[5] K. W. Mahmud, E. N. Duchon, Y. Kato, N. Kawashima, R. T. Scalettar, and N. Trivedi, Phys. Rev. B **84**, 054302 (2011).
[6] M. Greiner, O. Mandel, T. Esslinger, T. W. Hänsch, and I. Bloch, Nature **415**, 39 (2002).
[7] I. Bloch, J. Dalibard, and W. Zwerger, Rev. Mod. Phys. **80**, 885 (2008).
[8] W. Krauth, N. Trivedi, and D. Ceperley, Phys. Rev. Lett.

67, 2307 (1991).
[9] M. Wallin, E. S. Sørensen, S. M. Girvin, and A. P. Young, Phys. Rev. B **49**, 12115 (1994).
[10] J.-W. Lee, M.-C. Cha, and D. Kim, Phys. Rev. Lett. **87**, 247006 (2001).
[11] F. Pázmándi and G. T. Zimányi, Phys. Rev. B **57**, 5044 (1998).
[12] J. Wu and P. Phillips, Phys. Rev. B **78**, 014515 (2008).
[13] N. V. Prokof'ev and B. V. Svistunov, Phys. Rev. Lett. **92**, 015703 (2004).
[14] L. Pollet, N. V. Prokof'ev, B. V. Svistunov, and M. Troyer, Phys. Rev. Lett. **103**, 140402 (2009).
[15] V. Gurarie, L. Pollet, N. V. Prokof'ev, B. V. Svistunov, and M. Troyer, Phys. Rev. B **80**, 214519 (2009).
[16] F. Krüger, S. Hong, and P. Phillips, Phys. Rev. B **84**, 115118 (2011).
[17] F. Lin, E. S. Sørensen, and D. M. Ceperley, Phys. Rev. B **84**, 094507 (2011).
[18] S. G. Söyler, M. Kiselev, N. V. Prokofev, and B. V. Svistunov, arXiv:1107.5437.
[19] E. Altman, Y. Kafri, A. Polkovnikov, and G. Refael, Phys. Rev. Lett. **93**, 150402 (2004).
[20] E. Altman, Y. Kafri, A. Polkovnikov, and G. Refael, Phys. Rev. B **81**, 174528 (2010).
[21] B. Damski, J. Zakrzewski, L. Santos, P. Zoller, and M. Lewenstein, Phys. Rev. Lett. **91**, 080403 (2003).
[22] L. Fallani, J. E. Lye, V. Guarrera, C. Fort, and M. Inguscio, Phys. Rev. Lett. **98**, 130404 (2007).
[23] R. B. Griffiths, Phys. Rev. Lett. **23**, 17 (1969).
[24] B. M. McCoy, Phys. Rev. Lett. **23**, 383 (1969).
[25] T. Vojta, J. Phys. A, Math. Gen. **39**, R143 (2006).
[26] O. F. Syljuåsen and A. W. Sandvik, Phys. Rev. E **66**, 046701 (2002); O. F. Syljuåsen, Phys. Rev. E **67**, 046701 (2003).
[27] D. Stauffer and A. Aharony, *Introduction to Percolation Theory* (Taylor & Francis, Philadelphia, 1994).
[28] T. Roscilde and S. Haas, Phys. Rev. Lett. **99**, 047205 (2007).